



Therapeutics of female infertility

- -For women whose first menstruation is at very early age.
- -Fibroids in uterus, myoma.
- -Menses too early, too profuse, too long.
- -Menses return with slightest mental excitement.
- -Leucorrhoea – milky... (***Sepia***), acrid in young girls(***Caulophyllum***).
- Extremely sensitive to cold.
- -Craving for eggs and indigestible things.
- -Before menses - headache, colic, chilliness, leucorrhoea, tender breast (***Caulophyllum***)
- -Profuse sweating especially about the head.
- -Sourness of discharges.

■ **Calc carb**

- Menses-profuse, bright, early, discharge of blood between periods with sexual excitement. (***Ambra***).
- -Pain from sacrum to pubis, from below upwards shooting up the vagina.
- -Retained placenta from atony of uterus, promotes expulsions of moles or foreign bodies from uterus (***Canth***).
- -Haemorrhages with blood partly fluid and partly clotted.
- Tendency to miscarriage at 3rd month.
- Itching genitals, almost insatiable desire for coition.....

■ **Sabina**

- Heavy, sore, tender, ulcerated or prolapsed uterus.
- -Heavy dragging in pelvis.
-Fatigue or hysteria with prolapse or other uterine complaints
- -Sensation of heavy dragging of pelvis.
-Vulva aphthous.
- Great consciousness of the womb.
- Feels the uterus at all times, feels it move and shift with her movements.

- *Indurations, Cyst, tumours in uterus, ovaries, testicles.*
- Uterine tumours, cramping pain in uterus beginning with coldness in abdomen.
- -Ossified uterus. Uterus fills up the whole pelvis.
- -Chancres, ulcers, bubo, chronic uterine inflammation, prolapse, induration of ovary.
- *-Induration of one part of uterus, softening of another.*
- -Tumours of breast and uterus.
- -Atonic amenorrhoea, scanty and delayed menses.
- -Deficient sexual desire.
- -Sterility from ovarian dropsy.
- -Corrosive leucorrhoea with pustules on genitals.
Cramping pain in uterus.

■ **Aurum Muriaticum Natronatum**

- -Metrorrhagia at climacteric, profuse, for weeks.
- -Flow dark, clotted, offensive, in gushes, with malignant disease of uterus.
- -Intense menstrual colic, with drawing up of knees and terrible bearing down pain like labour, must press feet against support as in labor.
- -Sycotic sterility.
- -Intense pruritus of labia and vagina < by thinking of it, > rubbing and tap water bath, especially in non menstrual period.
- -Leucorrhoea-Thin and of fishy odour Sycotic taint - A/F suppressed gonorrhoea or inheritant gonorrhoea for women with chronic pelvic disorders.....
- **Medorrhinum**

- Menses-prolonged or vicarious with epistaxis.
- -Female breasts have erysipelatous appearance, red streaks, with thin ichorous discharge.
- -Suppuration of mammae, burning, watery, offensive discharges.
- -Drawing, forcing towards pelvis and rectum, as if menses would appear.
- For tall slender persons, with sanguine temperament.
Tubercular diathesis.

■ Phos

- Irregular menses of nearly every form, with pains going to the back.
 - Stasis and weakness with ptosis of viscera and weariness.
- -Dryness of vagina after menses.
 - Coition painful.
 - Aversion to coition. Nausea and irritability at thought of coitus.
- -A remedy for prolapse, violent stitches upwards in the vagina, lancinating pain from uterus to umbilicus.
- -Tendency to abortion from 5th to 7th month.....
- **Sepia**

- VAGINA SENSITIVE [***Berb, Kreos, Lyssin***], prevents coition (***Plat***).
- -SEVERE LEFT OVARIAN PAINS [***Crocus, Ustiligo***] < MENSES.
- -ABORTION in 3rd month.
- Sweats on uncovered parts
- -A/F-Vaccination, suppressed gonorrhoea.
- -It is a remedy for hydrogenoid or lymphatic constitution -susceptible to damp
- -Polyps on vagina and wart-like excrescences.....
- **Thuja**

- Uterus feels sore as if squeezed.
- -Hypertrophy of uterus.
- -Marked dysmenorrhoea.
- With an intense soreness, bruised feeling over the whole body. Soreness of abdominal walls and uterus.
- Injury to the deeper, internal organs.
- Left sided affections with a very sore; bruised feeling
- Also known as wound wort or Bruise wort.....
- **Bellis per.**

- -Itching and burning in vagina
- -Menses too late, short, scanty, difficult, thick, black, ACRID, MAKING PARTS SORE
- -During menses- headache, rush of blood to the head, nosebleed, pressure in pit of stomach.
- -MENORRHAGIA; HAS NOT BEEN WELL SINCE LAST MISCARRIAGE.
- Offensive, burning leucorrhoea.
- -Restlessness >perspiration. RESTLESSNESS BEFORE MENSES.
- -RED ORIFICES.
- **Sulphur**

- -Localised coldness or numbness; of single parts.
- -Menses too early, too profuse, long lasting; flow dark, black, clotted with painful bearing down spasms.
- -Violent cramping pains attended with numbness of the parts [***Cham, Acon, Kalm***].
- -Sleeps on back with knees drawn up spread wide apart; wants to uncover completely. Lascivious dreams.
- Nymphomania < in puerperal state. Eroticism, vaginismus with spasm, constriction.

■ Platina

- -Abdomen tenderness, in left inguinal region bearing down pain, extending down thigh.
- -Enlargement of the uterus.
- -Fibrous growths, subinvolution and prolapse.
- -Uterine tumors with bearing down sensation. Uterus enlarged, and patulous.
- -Fibroid with bearing down sensation.
- -Dysmenorrhoea.
- **Fraxinus Americanus**



■ THANK YOU